

the observation that 70 percent of the closely spaced double pulses are followed by a third pulse interpretable as a μ -decay while only 30 percent of all the sweeps show a second pulse.

(d) By extrapolation of the decay curve of Fig. 2 back to zero time, we conclude that about 1.5 π^+ -mesons were stopped in each gram of crystal per hour. A Kodak NTB nuclear emulsion placed near the crystal indicated that about 1.2 π^+ -mesons were stopped in each gram of emulsion per hour.

The circled points in Fig. 2 show the number of delayed pulses corrected for accidentals. By fitting an exponential curve to these points¹ one finds for the mean life of π^+ -mesons the value $\tau = (1.65 \pm 0.33) \times 10^{-8}$ sec. Within the rather large statistical errors involved, this value of the mean life agrees with the value $(1.97^{+0.14}_{-0.17}) \times 10^{-8}$ sec. obtained by Martinelli and Panofsky² on the decay of π^+ -mesons in flight, and with the value $(1.11^{+0.31}_{-0.22}) \times 10^{-8}$ sec. obtained by Richardson³ on the decay of π^- -mesons in flight.

We wish to thank the members of the M.I.T. synchrotron staff who have cooperated in making these measurements. Professors Bruno Rossi and Matthew Sands have contributed much encouragement and sound advice. We thank Mr. W. B. Smith who constructed and assisted in the testing of most of the equipment, and Mr. A. Grubman who grew the stilbene crystals.

† This work was supported in part by the joint program of the AEC and ONR.

* Physics Department and Laboratory for Nuclear Science and Engineering.

** Electrical Engineering Department and Laboratory for Nuclear Science and Engineering.

¹ R. Peierls, Proc. Roy. Soc. A149, 467 (1935).

² E. A. Martinelli and W. K. H. Panofsky, Phys. Rev. 77, 465 (1950).

³ J. R. Richardson, Phys. Rev. 74, 1720 (1948).

Superconductivity of Isotopes of Mercury*

C. A. REYNOLDS, B. SERIN, W. H. WRIGHT, AND L. B. NESBITT
Rutgers University, New Brunswick, New Jersey
March 24, 1950

THE superconducting transition temperatures of natural lead and the lead isotope obtained from the decay of uranium have been determined from resistance measurements by Kamerlingh Onnes and Tuyn¹ and by Justi.² In both instances, no detectable change in transition temperature between the natural element and the isotope was found.

In the measurements to be described, critical field *vs.* temperature curves for mercury samples enriched in various isotopes were determined by susceptibility measurements. The samples were obtained from the U. S. Atomic Energy Commission, and were prepared by electromagnetic separation. The average mass numbers for the enriched samples (samples 1, 2, 4) are shown in Table I; sample 3 is natural mercury.

The samples varying in mass from 50 to 100 mg, were sealed under helium in capillaries about 0.5 mm i.d., giving samples about 3 cm long. A coil of No. 41 copper wire was wound around each capillary. The samples were placed in a Dewar vessel of liquid helium, and formed the secondary of a mutual inductance with a solenoid placed outside the shield Dewar. The solenoid was excited by a 1000 c/sec. oscillator, and produced a magnetic field at the sample of less than 0.1 oersted. The signal picked up by the sample coil was amplified and detected. The signal was balanced out with the sample superconducting, and then the detected signal

TABLE I. Transition temperatures.

Sample	Average mass number	T_0 (°K)
1	203.4	4.126
2	202.0	4.143
3	200.7	4.150
4	199.7	4.161




FIG. 1. The critical magnetic field as a function of the absolute temperature.

was observed as a function of the magnetic field of a large Helmholtz coil surrounding the Dewars. The earth's magnetic field was canceled to 0.3 percent. When the magnetic field reached the neighborhood of the critical value at any temperature, there was a rapid increase in detected signal. The critical field was taken to be the average value in the transition interval. Figure 1 shows the critical field curves for the four samples. It is to be noted that there is a systematic decrease of transition temperature with increasing mass. The transition temperatures (in zero magnetic field), T_0 , of the samples are given in Table I. The slopes of these curves at zero field agree within 1.0 percent; the magnitude of the slope is -204 ± 2 oersteds/°K.

Figure 2 is a plot of transition temperature *vs.* average mass




FIG. 2. The transition temperature *vs.* the average mass numbers of the isotopic mixtures.

number. The values obtained by Maxwell³ for Hg¹⁹⁸ and natural Hg are included. The slope of the line in Fig. 2 is 0.009 °K/(mass number).

* This work was supported by the ONR, by the Research Corporation, by the Rutgers University Research Council, and by the Radio Corporation of America.

¹ H. Kamerlingh Onnes and W. Tuyn, Leiden Comm. No. 160b (1922).

² E. Justi, Physik. Zeits. 42, 325 (1941).

³ E. Maxwell, Phys. Rev. 78, 477 (1950).