

Method and Means for Hypergeometric Function Calculation on an Array Processor

Author(s): **IBM TDB
Grant, PM**

IP.com number: **IPCOM000054707D**

Original Publication Date: **March 1, 1980**

Original Disclosure Information: **TDB 03-80 p4699-4701**

IP.com Electronic Publication: **February 13, 2005**

IP.com, Inc. is the world's leader in defensive publications. The largest and most innovative companies publish their technical disclosures into the IP.com Prior Art Database. Disclosures can be published in any language, and they are searchable in those languages online. Unique identifiers indicate documents containing chemical structures as well as publications open for comment in the IP Discussion Forum. Disclosures are published every day online and also appear in the printed IP.com Journal. The IP.com Prior Art Database is freely available to search by patent examiners throughout the world.

Client may copy any content obtained through the site for Client's individual, noncommercial internal use only. Client agrees not to otherwise copy, change, upload, transmit, sell, publish, commercially exploit, modify, create derivative works or distribute any content available through the site.

Note: This is a pdf rendering of the actual disclosure. To access the notarized disclosure package containing an exact copy of the publication in its original format as well as any attached files, please download the full document from the IP.com Prior Art Database at: <http://www.ip.com/pubview/IPCOM000054707D>

Method and Means for Hypergeometric Function Calculation on an Array Processor

This article describes a method for efficient calculation in the hardware of an array processor, of those hypergeometric functions which are defined by successive differentiation of a seed function. Examples are the Legendre functions, defined by:

$$P(l)(z) = (1-z/2)^{l/2} \frac{d^l P(l)(z)}{dz^l}$$

where $P(z)$ is the l -th Legendre polynomial; the associated Laguerre functions, defined by:

$$L_s(r)(\rho) = \frac{d^s}{d\rho^s} L(r)(\rho)$$

where $L(P)$ is the P -th Laguerre polynomial; and the spherical Bessel functions, defined by:

$$j(n)(z) = z^{-n} \frac{d^n}{dz^n} \frac{\sin z}{z}$$

The attribute which permits the utilization of an array processor in the calculation of such functions resides in the fact that the seed function can be usually represented as a polynomial whose general term is simply:

$$a(n) z^n$$


with general derivative

See Original.

Thus, if in the calculation of the seed function polynomial either by recursion as in the case of Legendre and Laguerre polynomials, or by economization as for spherical Bessel functions, one forms an appropriate array of coefficients and exponents, the calculation of the required hypergeometric function becomes a trivial shift and multiply operation of this array in hardware capable of parallel array operations. Often the seed function itself can be calculated via such array manipulations, as, for example, when we define the Legendre polynomials by Rodriguez' formula: See Original.

In function ALF, lines 10-12 compute the coefficients and exponents of the seed Legendre polynomial by recursion. Lines 13-14 compute the required value of the associated Legendre function by the array manipulation algorithm discussed in this article.

The figure diagrammatically depicts the parallel processor architecture necessary for calculating hypergeometric function through application of Eqs. (4) and (5). The a_n , Z_n , and n arrays are assumed to be previously loaded as is also the m -counter. The result will be accumulated in the n array. The n array is provided with a cumulative multiplier, that is, the product developed in the i th element is the product of $i, i-1, i-2, \dots, i-m+1$. The numbers 1, 2, and 3 emanating from the clock denote the timing sequence.



TIMING: 1 BEFORE 2 BEFORE 3

$$P_{\ell m}(z) = (1-z^2)^{m/2} \frac{d^m P_{\ell}(z)}{dz^m} .$$

$$L_r^s(\rho) = \frac{d^s}{d\rho^s} L_r(\rho), \quad j_n(z) = z^n \left(-\frac{1}{z} \frac{d}{dz} \right)^n \frac{\sin z}{z} . \quad \frac{d^m}{dz^m} a_n z^n = a_n \left(\frac{n!}{(n-m)!} \right) z^{n-m} .$$

$$P_{\ell}(z) = \frac{1}{2^{\ell} \ell!} \frac{d^{\ell}}{dz^{\ell}} (z^2 - 1)^{\ell} .$$